

A better large N expansion for chiral Yukawa models.*

George Bathas^a and Herbert Neuberger^{a †}

^aDepartment of Physics and Astronomy
Rutgers University, Piscataway, NJ 08855, U.S.A.

We consider the most general renormalizable chiral Yukawa model with $SU(3)_{\text{color}}$ replaced by $SU(N_c)$, $SU(2)_L$ replaced by $SU(N_w)$ and $U(1)_Y$ replaced by $U(1)^{N_w-1}$ in the limit $N_c \rightarrow \infty$, $N_w \rightarrow \infty$ with the ratio $\rho = \sqrt{\frac{N_w}{N_c}} \neq 0, \infty$ held fixed. Since for $N_w \geq 3$ only one renormalizable Yukawa coupling per family exists and there is no mixing between families the limit is appropriate for the description of the effects of a heavy top quark when all the other fermions are taken to be massless. The large $N = \sqrt{N_c N_w}$ expansion is expected to be no worse quantitatively in this model than in the purely scalar case and the $N = \infty$ limit is soluble even when the model is regularized non-perturbatively. A rough estimate of the triviality bound on the Yukawa coupling is equivalent to $m_t \leq 1 \text{ TeV}$.

We wish to study non-perturbatively a model that is reasonably close to what the minimal standard model would look like if the top quark Yukawa coupling and the Higgs selfcoupling were both large while all other couplings were kept at their known small values and could therefore be neglected in a first approximation.

We propose to do this using an $\frac{1}{N}$ expansion on a model defined non-perturbatively with a Pauli Villars type of regularization[1]. The action is:

$$\begin{aligned} -S = & \int_x \bar{\psi}_{-}^{ia} \not{\partial}_+ \psi_{+}^{ia} + \bar{\chi}_{+}^a h_{\chi} (-\partial^2) \not{\partial}_- \chi_{-}^a \\ & + \phi^{i*} h_{\phi} (-\partial^2) \partial^2 \phi^i - m_0^2 \phi^{i*} \phi^i - \frac{\lambda_0}{4N_w} (\phi^{i*} \phi^i)^2 \\ & + \frac{g_0}{\sqrt{N}} [\bar{\psi}_{-}^{ia} \chi_{-}^a \phi^i + \bar{\chi}_{+}^a \psi_{+}^{ia} \phi^i]; \quad g_0 > 0 \end{aligned} \quad (1)$$

where

$$\begin{aligned} N = & \sqrt{N_w N_c}, \quad \not{\partial}_+ = \sigma_{\mu} \partial_{\mu}, \quad \not{\partial}_- = \bar{\sigma}_{\mu} \partial_{\mu} \\ \bar{\sigma}_{1,2,3} = & \sigma_{1,2,3}, \quad \bar{\sigma}_4 = -\sigma_4 = -i \\ i = & 1, \dots, N_w \quad a = 1, \dots, N_c \end{aligned} \quad (2)$$

The $\sigma_{1,2,3}$ are Pauli matrices. Ultraviolet divergences can be eliminated by just regularizing the χ and ϕ propagators because there are no loops

*This research was supported in part by the DOE under grant # DE-FG05-90ER40559.

[†]Speaker at the Conference.

consisting of only ψ propagators. This is achieved with $h_{\chi}(p^2) = h_{\phi}(p^2) = 1 + (p^2/\Lambda^2)^n$ with $n \geq 3$.

ψ represents the top-bottom doublet, χ the right handed top and ϕ the Higgs doublet. The index i generalizes the weak isospin index and the index a labels colors. ψ_{+} is a Euclidean left handed Weyl spinor, χ_{-} a right handed one and ϕ a complex scalar. $\bar{\psi}_{-}$ and $\bar{\chi}_{+}$ are independent fields of handedness opposite to their unbarred counterparts as usual in the Euclidean functional integral formulation.

The large N limit is taken with $\bar{\rho} \equiv \sqrt{\frac{N_w}{2N_c}}$ held fixed and is dominated by planar diagrams. Renormalizability allows us to get away with not having to introduce four Fermion interactions and with them an insoluble planar diagram problem, at no cost to generality.¹ Thus the model is manageable being similar diagrammatically to two dimensional QCD with matter at large N_c . The large N limit can be found by looking at the structure of planar Feynman graphs: essentially one sums over all ‘‘cactuses of bubbles’’ and all ‘‘rainbows’’. For future work it is somewhat preferable to use functional integral manipulations to the same end.

As an example of what can be done in this scheme we present a very simple calculation: We compute a β -function associated with the

¹Except for possible cutoff effects.

Yukawa coupling constant defined in some simple manner and ignoring cutoff effects. This can be done with relative ease because even at infinite order in the couplings the flow of this Yukawa coupling is unaffected by λ_0 , similarly to the finite N form at one loop order. From this calculation a rough estimate for the “triviality” bound on the top mass can be obtained.²

To find β functions it suffices to solve the model at criticality when all masses vanish. We start in the symmetric phase and pick a mass independent renormalization scheme to make the zero mass limit smooth.

We introduce bilocal fields,

$$\begin{aligned} S_{\alpha,\beta}(x,y) &= \frac{1}{N_c} \bar{\chi}_{+\alpha}^a(x) \chi_{-\beta}^a(y) \\ K(x,y) &= \frac{1}{N_w} \phi^{i*}(x) \phi^i(y) \end{aligned} \quad (3)$$

where α, β are space-time spinor indices, into the functional integral by writing representations for the appropriate δ -functions with the help of two auxiliary bilocal fields $\lambda_{\alpha,\beta}(x,y)$ and $\mu(x,y)$. We can integrate out now all the original fields, including the zero mode of the scalar field because we assumed that we are in the symmetric phase. This makes the N and $\bar{\rho}$ dependence explicit and, when $N \rightarrow \infty$, the integral over S, λ, K and μ is dominated by a saddle point which has to satisfy the following equations:

$$\begin{aligned} K(x,y) &= \{[-h_\phi(-\partial^2)\partial^2 + m_0^2 + \mu]^{-1}\}_{y,x} \\ S_{\alpha,\beta}(x,y) &= \{[h_\chi(-\partial^2)\bar{\phi} + \lambda]^{-1}\}_{y\beta,x\alpha} \\ \lambda_{\alpha,\beta}(x,y) &= -g^2 \bar{\rho} \sqrt{2} \{\bar{\phi}_+^{-1}\}_{x\alpha,y\beta} K(x,y) \\ \mu(x,y) &= \frac{g^2}{\bar{\rho} \sqrt{2}} \sum_{\alpha,\beta} \{\bar{\phi}_+^{-1}\}_{x\alpha,y\beta} S_{\alpha,\beta}(x,y) \\ &\quad + \frac{\lambda_0}{2} K(x,x) \delta_{x,y} \end{aligned} \quad (4)$$

We go to Fourier space and use translational and rotational invariance to write

$$\tilde{\lambda}_{\alpha,\beta}(p) = ip_\mu \bar{\sigma}_\mu \hat{\lambda}(p^2) \quad \tilde{\mu}(p) = \hat{\mu}(p^2) \quad (5)$$

Let us introduce Z_χ and Z_ϕ , wave function renormalization constants, and new functions $f(p^2) =$

²In practice this bound might be somewhat uninteresting because it is likely to be higher than the vacuum stability bound for reasonable Higgs masses.

$Z_\chi[\hat{\lambda}(p^2) + h_\chi(p^2)]$ and $g(p^2) = Z_\phi[p^2 h_\phi(p^2) + m_0^2 + \hat{\mu}(p^2)]$. Since at the planar level there are no vertex corrections to the Yukawa coupling³ it makes sense to define a renormalized Yukawa coupling by $g_R^2 = Z_\phi Z_\chi g_0^2$, ($\alpha \equiv \frac{g_R^2}{16\pi^2}$). We also set $\lambda'_0 = \lambda_0 Z_\phi^2$ but, unlike α , do not expect λ'_0 to stay finite when $\Lambda \rightarrow \infty$. Z_χ and Z_ϕ are defined in terms of the bare couplings by $f(\mu^2) = \frac{g(\mu^2)}{\mu^2} = 1$.

After performing the angular part of the momentum integrals one gets when the products in the saddle point equations for x, y dependent quantities are converted into convolutions in momentum space one finds:

$$\begin{aligned} f(p^2) &= Z_\chi h_\chi(p^2) + \sqrt{2}\alpha \bar{\rho} \left[\frac{1}{2} \int_{p^2}^\infty \frac{dk^2}{g(k^2)} \right. \\ &\quad \left. + \frac{1}{p^2} \int_0^{p^2} \frac{k^2 dk^2}{g(k^2)} - \frac{1}{2p^4} \int_0^{p^2} \frac{k^4 dk^2}{g(k^2)} \right] \\ g(p^2) &= Z_\phi p^2 h_\phi(p^2) + \frac{\sqrt{2}\alpha}{\bar{\rho}} \left[\frac{p^2}{2} \int_{p^2}^\infty \frac{dk^2}{k^2 f(k^2)} \right. \\ &\quad \left. + \int_0^{p^2} \frac{dk^2}{f(k^2)} - \frac{1}{2p^2} \int_0^{p^2} \frac{k^2 dk^2}{f(k^2)} \right] + m^2 \\ m^2 &= Z_\phi m_0^2 + \frac{\lambda'_0}{32\pi^2} \int_0^\infty \frac{k^2 dk^2}{g(k^2)} \\ &\quad - \frac{\sqrt{2}\alpha}{\bar{\rho}} \int_0^\infty \frac{dk^2}{f(k^2)} \end{aligned} \quad (6)$$

The massless case is obtained by adjusting m_0^2 so that $m^2 = 0$ and then it makes sense to define $h(p^2) = \frac{g(p^2)}{p^2}$ leading to a set of equations of a more symmetric appearance with an interesting bosonic-fermionic symmetry at $\bar{\rho} = 1$.

Rescaling the functions and the momentum variable by

$$f(u\mu^2) = \sqrt{2\alpha} \bar{\phi}(u) \quad h(u\mu^2) = \sqrt{\alpha} \bar{\psi}(u) \quad (7)$$

we obtain:

$$\begin{aligned} \bar{\psi}(u) &= \frac{1}{\sqrt{\alpha}} + \frac{Z_\phi}{\sqrt{\alpha}} [h_\phi(u\mu^2) - h_\phi(\mu^2)] \\ &\quad + \frac{1}{\bar{\rho}} \int_u^1 \frac{dv}{v} \int_0^1 \frac{(1-x)dx}{\bar{\phi}(xv)} \\ \bar{\phi}(u) &= \frac{1}{2\sqrt{\alpha}} + \frac{Z_\chi}{2\sqrt{\alpha}} [h_\chi(u\mu^2) - h_\chi(\mu^2)] \\ &\quad + \bar{\rho} \int_u^1 \frac{dv}{v} \int_0^1 \frac{(1-x)dx}{\bar{\psi}(xv)} \end{aligned} \quad (8)$$

³This is also true at one loop order at finite N .

It is clear that when n is large enough h and f are dominated by the $h_{\phi,\chi}$ functions in the ultraviolet and therefore the large cutoff behavior of the wave function renormalization constants is such that in the infinite cutoff limit, at u and μ^2 fixed, Z_ϕ and Z_χ simply disappear from the above equation, leaving us with

$$\begin{aligned}\bar{\psi}(u) &= \frac{1}{\sqrt{\alpha}} + \frac{1}{\bar{\rho}} \int_u^1 \frac{dv}{v} \int_0^1 \frac{(1-x)dx}{\bar{\phi}(xv)} \\ \bar{\phi}(u) &= \frac{1}{2\sqrt{\alpha}} + \bar{\rho} \int_u^1 \frac{dv}{v} \int_0^1 \frac{(1-x)dx}{\bar{\psi}(xv)}\end{aligned}\quad (9)$$

On physical grounds it is obvious that we wish that $\bar{\psi}$ and $\bar{\phi}$ be positive. The equations above show that if this is true in some interval $(0, u_*)$ then the functions will be monotonically decreasing there. However, the equation do not admit an asymptotic behavior as $u \rightarrow \infty$ with both functions approaching non-negative limits so the positivity requirement must get violated somewhere in the ultraviolet.⁴ Starting at the scale where the violation first occurs cutoff effects cannot be neglected any more. This “unphysical” scale is the usual Landau pole, this time appearing in a nonperturbative approximation.

In the infrared there is no Landau pole problem and cutoff effects are indeed negligible. To investigate the behavior there we set $x = -\log(u)$ and $\hat{\psi}(x) = \bar{\psi}(u)$, $\hat{\phi}(x) = \bar{\phi}(u)$ and derive:

$$\begin{aligned}\mathcal{D}\hat{\psi} &= \frac{1}{\bar{\rho}\hat{\phi}}, & \mathcal{D}\hat{\phi} &= \frac{\bar{\rho}}{\hat{\psi}} \\ \mathcal{D} &\equiv \frac{d}{dx} \left(\frac{d}{dx} - 1 \right) \left(\frac{d}{dx} - 2 \right)\end{aligned}\quad (10)$$

These equations allow us to extract the infrared behavior of the solution of the integral equations. We ended up with only differential equations (rather than integral) reflecting that according to the Renormalization Group one needs only very limited information at a given scale in order to derive the behavior at a scale close by; therefore a differential equation must show up eventually, its order less the number of asymptotic conditions determining the amount of information at a given scale that is needed to determine

⁴A zero of $\bar{\psi}$ or $\bar{\phi}$ at a positive u corresponds to a pole in a two-point function at an Euclidean momentum; the “particle” associated with this pole would be tachyonic.

the behavior at the next scale. Our choice of regularization was made so that even at finite cutoff $\hat{\phi}$ and $\hat{\psi}$ obey a set of purely differential equations of a structure similar to 10.

Since the fixed point governing the infrared behavior is the free field fixed point the form of the solutions in the infrared simply embodies the two anomalous dimensions associated with the χ and ϕ fields when expanded in α around $\alpha = 0$. These anomalous dimensions as a function of the coupling also determine the β function and from the asymptotic series of $\psi(x)$ and $\phi(x)$ at $x \rightarrow \infty$ we can get the contributions to the $\beta(\alpha)$ function ordered in the number of loops.

The differential equations lead to:

$$\begin{aligned}\hat{\psi} &\sim \psi_0 x^b [1 + c \frac{\log x}{x} + c' \frac{1}{x} + \dots] \\ \hat{\phi} &\sim \phi_0 x^a [1 + d \frac{\log x}{x} + d' \frac{1}{x} + \dots]\end{aligned}\quad (11)$$

where

$$\begin{aligned}a &= b\bar{\rho}^2 = \frac{\bar{\rho}^2}{1 + \bar{\rho}^2}, & d &= c\bar{\rho}^2 = -\frac{3\bar{\rho}^4}{(1 + \bar{\rho}^2)^3} \\ \frac{d'}{\bar{\rho}^2} - c' &= \frac{3(1 - \bar{\rho}^2)}{2(1 + \bar{\rho}^2)^2}; & \psi_0\phi_0 &= \frac{\bar{\rho} + \bar{\rho}^{-1}}{2}\end{aligned}\quad (12)$$

Only the product $\psi_0\phi_0$ and a particular linear combination of d' and c' get determined because the equations are invariant under a field rescaling $\hat{\psi} \rightarrow A^{-1}\hat{\psi}$, $\hat{\phi} \rightarrow A\hat{\phi}$ reflecting a change in the finite parts of the wave function renormalization constants and a shift $x \rightarrow x + x_0$ reflecting a change in μ^2 . The field-rescaling invariance disappears in the product $r(u) \equiv \sqrt{2}\bar{\psi}(u)\bar{\phi}(u)$ and this combination is indeed special because it satisfies a Renormalization Group equation in which only the β -function appears (the sum of the two anomalous dimensions rather than each individual one):

$$\left[-\frac{\partial}{\partial t} + \beta(\alpha) \frac{\partial}{\partial \alpha} \right] r(e^{2t}, \alpha) = 0 \quad (13)$$

This equation is exact in our limit. From it we derive another exact relation between r and β :

$$\beta(\alpha(u\mu^2)) = -2[\alpha(u\mu^2)]^2 \frac{dr(u)}{d(\log u)} \Big|_{r(u)=\frac{1}{\alpha(u\mu^2)}} \quad (14)$$




Figure 1. The nonperturbatively determined β -function for the Yukawa coupling at $N_c, N_w \rightarrow \infty$ with $\frac{N_c}{N_w} = \frac{3}{2}$. At tree level the top quark mass is given by $m_t = (4\pi^2\alpha/3)^{\frac{1}{2}} 246 [GeV]$: $\beta(\alpha) = \frac{d\alpha}{dt}$ where t is the logarithm of the energy scale.

Using now the asymptotic expansion of the exact solution in the infrared we get:

$$\beta(\alpha) = \alpha^2[\sqrt{2}(\bar{\rho} + \bar{\rho}^{-1}) - 3\alpha + O(\alpha^2)] \quad (15)$$

The one loop result when expressed in a conventional variable, $y(t)$ (in the standard model the top mass at tree level is $m_t = y(\sim 0) 246 [GeV]$), is, with the true value $\bar{\rho} = \frac{1}{\sqrt{3}}$, $\frac{dy^2}{dt} = \frac{y^4}{\pi^2}$ which is close to the finite N result, $\frac{dy^2}{dt} = \frac{9y^4}{8\pi^2}$. The difference comes from the wave function renormalization of the left handed ψ field which is suppressed at leading order in $\frac{1}{N}$. For arbitrary N_c and N_w we would have obtained at one loop order $\frac{dy^2}{dt} = \frac{(2N_c + N_w + 1)y^4}{8\pi^2}$ and we see that the relative magnitude of the $1/N$ correction is much smaller than in the scalar case[2].

To get the full β -function at infinite N we solve for $\hat{\psi}$ and $\hat{\phi}$ numerically by iterating an integral form of the equations 6 but with the normaliza-

tion conditions at μ^2 incorporated; this avoids the appearance of the double integrals in 8 and 9.

The result for β is shown in the figure and one again sees explicit evidence for the nonperturbative existence of a Landau pole. We compare the Landau pole energy obtained from the exact result to the one loop estimate. For example, with the high value of $\alpha(\mu^2) = 1.365$ chosen in the figure we obtain $\Lambda_{Landau}^2 = 2\mu^2$ while at one loop $\Lambda_{Landau}^2 = 1.565\mu^2$. A top quark corresponding to such a strong coupling would have a mass of $m_t = 4\pi\sqrt{\frac{\alpha}{12}} 246 [GeV] \approx 1043 GeV$. We see that “triviality bounds” on the top will likely come out close to the perturbative unitarity bounds of Chanowitz et. al.[3] in contrast to Einhorn and Goldberg[4] who obtained $5600 GeV$.

From our non-perturbative result in the figure one sees that the one and two loop errors in β are about comparable for $\alpha \approx 0.4$ (roughly of order 20 percent) and that using the one loop results all the way out to infinity is not very harmful, but the two loop result would be completely misleading if extrapolated beyond its region of validity, to $\alpha \approx 1$. A top mass of $200 GeV$ would correspond to $\alpha \approx 0.05$, well within the perturbative domain.

For the future we plan the following steps:

- Compute the β -function associated with λ at infinite cutoff in the massless case.
- Reintroduce the finite cutoff in the above computation to see the set in and magnitude of cutoff scaling violations.
- Redo both the infinite cutoff and the finite cutoff computation away from the critical line, i.e. when thresholds are present. This has to be done in both the broken and the symmetric phases.
- Compute the effective potential at infinite cutoff limit first and then at finite cutoff and analyze the issue of “vacuum instability” in this non-perturbative setting.
- Study bound state problems.

REFERENCES

1. G. Bathas, H. Neuberger, **Phys. Lett. B**, to appear.
2. R. Dashen and H. Neuberger, **Phys. Rev. Lett.** **50** (1983) 1897.
3. M. S. Chanowitz, M. A. Furman and I. Hinch-

- liffe, **Nucl. Phys. B153** (1979) 403.
4. M. B. Einhorn and G. Goldberg, **Phys. Rev. Lett. 57** (1986) 2115.